

REVIEW

Advantages of an antagonist: bicuculline and other GABA antagonists

Graham AR Johnston

Pharmacology, School of Medical Sciences, The University of Sydney, NSW, Australia

Correspondence

Emeritus Professor Graham AR Johnston, Pharmacology, School of Medical Sciences, The University of Sydney, Sydney NSW 2006 Australia. E-mail: grahamj@mail.usyd.edu.au

Keywords

GABA receptors; antagonists; bicuculline; gabazine; picrotoxin

Received

11 December 2012

Revised

21 January 2013

Accepted

31 January 2013

The convulsant alkaloid bicuculline continues to be investigated more than 40 years after the first publication of its action as an antagonist of receptors for the inhibitory neurotransmitter GABA. This historical perspective highlights key aspects of the discovery of bicuculline as a GABA antagonist and the sustained interest in this and other GABA antagonists. The exciting advances in the molecular biology, pharmacology and physiology of GABA receptors provide a continuing stimulus for the discovery of new antagonists with increasing selectivity for the myriad of GABA receptor subclasses. Interesting GABA antagonists not structurally related to bicuculline include gabazine, salicylidene salicylhydrazide, RU5135 and 4-(3-biphenyl-5-(4-piperidyl)-3-isoxazole. Bicuculline became the benchmark antagonist for what became known as GABA_A receptors, but not all ionotropic GABA receptors are susceptible to bicuculline. In addition, not all GABA_A receptor antagonists are convulsants. Thus there are still surprises in store as the study of GABA receptors evolves.

Abbreviations

gabazine, SR 95531, 4-[6-imino-3-(4-methoxyphenyl)pyridazin-1-yl] butanoic acid; RU5135, 3 α -hydroxy-16-imino-17-aza-5 β -androstane-11-one; 4-PIOL, 5-(4-piperidyl)-3-isoxazolol

Introduction

The editorial accompanying our 1970 Nature paper on the GABA receptor antagonist action of the convulsant alkaloid bicuculline (Curtis *et al.*, 1970b) was entitled 'Advantages of an Antagonist' (Editorial, 1970). This was a very enthusiastic editorial beginning with 'Students of nervous transmission should be stimulated to a new burst of enthusiasm' and ending with 'With this tool it should now be possible to map fairly rapidly the distribution of GABA-inhibitory synapses in the CNS, and to determine whether they are as numerous and widely distributed as the relatively high GABA content of the tissue would suggest.' Interest in the GABA antagonist action of bicuculline has been remarkably sustained with an average of more than 120 publications per year since 1970 containing the terms 'GABA' and 'bicuculline' in their title or abstract according to the American Chemical Society's Chemical Abstracts and related databases as accessed via SciFinder. Only picrotoxin/picrotoxinin, averaging 70 publications per year since 1970, and gabazine, averaging 19 publications per year since it was introduced in 1986, rival bicuculline as relatively widely used GABA_A receptor antagonists.

Thanks to the pioneering work of Gene Roberts and others by 1970 (Roberts *et al.*, 1970) there had been much speculation about the possible function of GABA as an inhibitory transmitter in the CNS based on its distribution, metabolism and release from nervous tissue and the observation that GABA hyperpolarized neurons in a manner similar to synaptic inhibition. It was known that the convulsant alkaloid strychnine was able to antagonize synaptic inhibition in the spinal cord without influencing the inhibitory action of GABA. This puzzling observation was explained by the important findings by Werman *et al.*, (1967) that the simple amino acid glycine was likely to be an inhibitory transmitter in the spinal cord and our subsequent finding that strychnine antagonized the inhibitory action of glycine but not that of GABA (Curtis *et al.*, 1967). This work was a good example of the advantages of a selective antagonist.

The search was then on for a similarly selective antagonist of the inhibitory action of GABA that did not influence that of glycine. An obvious place to start was known convulsants. In the late 1960s, there were no readily available electronic databases; one had to hunt through paper versions of Chemical Abstracts, Index Medicus and the like. The Merck Index

actually proved to be the most useful in providing leads that could then be followed up in the original literature. A pattern emerged that many alkaloids containing the tetrahydroisoquinoline nucleus were convulsants. As we acquired these alkaloids from our chemist colleagues and chemical suppliers and tested them *in vivo* on neurons in the cat spinal cord, it became clear that most were glycine antagonists like strychnine without effect on the action of GABA.

Bicuculline as a GABA receptor antagonist

On 17 March 1970 (St Patrick's Day), David Curtis, Arthur Duggan, Dominik Felix and I, first tested bicuculline in the spinal cord of a cat under pentobarbitone anaesthesia. By 19 May, we had submitted a manuscript, 'GABA, bicuculline and central inhibition', to *Nature* that was published on 27 June (Curtis *et al.*, 1970b). An example of the selective antagonist action of bicuculline on the action of GABA is shown in Figure 1 where bicuculline reduced the effect of GABA without influencing the action of glycine.

The *Nature* issue of 14 November contained three papers on bicuculline and GABA: Hugh McLennan's findings on the antagonist action of bicuculline on GABA responses in crayfish (McLennan, 1970), an article by Godfraind *et al.* (Godfraind *et al.*, 1970) on the 'Doubtful value of bicuculline as a specific antagonist of GABA' and our response to the latter 'Bicuculline and central GABA receptors' (Curtis *et al.*, 1970a). Godfraind *et al.* concluded 'Whatever the mechanism of its convulsant action, we find it difficult to see how bicuculline could be useful for critical tests of GABA-mediated inhibition'. Their negative findings have never been satisfactorily explained but caused us to undertake elemental and spectroscopic analyses to confirm the chemical identity of the substance we had tested (Curtis *et al.*, 1970a). It seems unlikely that the 32 neurons that they examined in four cats had predominantly bicuculline-insensitive GABA receptors,

later classified as GABA_B receptors. More likely, the chemical instability of aqueous solutions of bicuculline was the problem. Their microelectrodes were filled with their bicuculline solutions 2–3 days before testing. While they were able to clearly show the convulsant properties of these test solutions taken from their microelectrodes, the potency appeared to be much reduced; they quoted pronounced convulsions being observed on systemic administration of bicuculline (5 μ mole). This works out at around 1.8 mg kg⁻¹ whereas many investigators have reported convulsions at 0.1 mg kg⁻¹.

Chemical instability of bicuculline

Many years earlier, Welch and Henderson (Welch and Henderson, 1934) had noted the slow conversion of bicuculline in solution to bicucine, a much less active convulsant (Figure 2). The chemical instability of bicuculline was further investigated by Olsen *et al.*, (1975). The conversion of bicuculline to bicucine involved the opening of a lactone ring and this could be readily monitored by UV spectroscopy. This conversion took place most readily at physiological pH. Bicuculline was relatively more stable at acidic pH as used in microelectrode studies. Bicucine was slowly converted to bicuculline at acidic pH. Quaternary salts of bicuculline, for example, bicuculline methiodide ('N-methyl bicuculline') (Pong and Graham, 1972) or methochloride (Johnston *et al.*, 1972), are much more stable than bicuculline, more water soluble and of similar potency as GABA antagonists but they do not appear to cross the blood brain barrier on systemic administration. Injected intracisternally, bicuculline methiodide was a more potent convulsant than bicuculline (Pong and Graham, 1972). While the quaternary salts are much easier to use than bicuculline, which exists as a protonated cation at physiological pH, care should be taken when using them, as they appear to be much less selective than bicuculline with respect to interactions with other systems (Seutin and Johnson, 1999). Bicuculline and its quaternary salts are agents with different pharmacological effects and sometimes it is not clear as to which agent was used in a particular study;





Figure 1

Differential effect of bicuculline, 40 nA ejection current, on the inhibition of a Renshaw cell by glycine (5 nA) and GABA (8 nA) in the spinal cord of a cat anaesthetized by pentobarbitone. Drugs were administered extracellularly from a multibarreled micropipette via electrophoresis. The firing frequency of the cell was maintained at around 60 spikes per second by the continuous electrophoretic ejection of DL-homocysteic acid (15 nA).

**Figure 2**

Structure of the protonated forms of the active GABA receptor antagonist bicuculline and the inactive bicucine that may be interconverted dependent on pH with bicucine favoured at physiological pH.


sometimes a study appears to be about bicuculline whereas in fact it involves a quaternary salt (Ueno *et al.*, 1997).

Earlier work on bicuculline

After publication of our work on bicuculline as a GABA receptor antagonist, we learnt of studies in the 1960s by Dr Kang Tsou in the Department of Pharmacology of the Shanghai Institute of Materia Medica. He had found that bicuculline was a potent convulsant when applied to the cerebral cortex and that it also reduced strychnine-insensitive inhibition of spinal reflexes. He presented these results at a Congress of the Chinese Society for Physiological Sciences in 1964 and submitted a full-length paper to *Acta Physiologica Sinica* in 1965 that was never published due to the suspension of the publication of all Chinese scientific journals during the Cultural Revolution. He was eventually able to publish on the CNS effects of bicuculline in 1976 in a new Chinese language journal, *Diseases of the Nervous System*. I was delighted to meet Dr Tsou in Shanghai in 1984 during a visit of Australian pharmacologists to China (Johnston, 1985) and to discuss his work on bicuculline (Figure 3). I saw some of his original records that clearly supported his original claims (Tsou, 1987). An excellent article on his career 'Lost in the cultural revolution' was published in 1994 (Hoffman, 1994).

GABA as a neurotransmitter

The advantage of bicuculline as a GABA receptor antagonist quickly became apparent as we started to investigate the influence of bicuculline on strychnine-insensitive synaptic inhibition. In our initial 1970 *Nature* paper (Curtis *et al.*, 1970b), we showed that bicuculline reduces strychnine-insensitive inhibition of pyramidal cells in the cerebral cortex and Purkinje cells in the cerebellum to 'provide for the first time decisive pharmacological evidence that, of the amino

**Figure 3**

Dr Kang Tsou in Shanghai in 1984 with his original records showing bicuculline influencing synaptic inhibition.

acids present in the mammalian nervous system, GABA is most likely to be the actual transmitter at these inhibitory synapses. The use of bicuculline and related substances will be of considerable assistance in studying other synapses for which GABA has been proposed as a possible transmitter.' We subsequently showed that bicuculline reduced presynaptic inhibition of spinal reflexes and the accompanying primary afferent depolarization (Curtis *et al.*, 1971).

GABA receptor subtypes

The division of GABA receptors into GABA_A and GABA_B subtypes dates from the studies of David Hill and Norman Bowery in 1981 on the binding of baclofen and GABA to rat brain membranes (Hill and Bowery, 1981). They described a receptor that 'differs from the classical GABA site as it is unaffected by recognized GABA antagonists such as bicuculline'. They went on to state 'We propose to designate the classical site as the GABA_A and the novel site as the GABA_B receptor.' GABA_A receptors were ionotropic receptors that gate chloride channels, while GABA_B receptors were G-protein coupled receptors. These receptors could be differentiated on the basis of agonist and antagonist selectivity. GABA_A receptors were antagonized by bicuculline and insensitive to baclofen, whereas GABA_B receptors were activated by baclofen and insensitive to bicuculline. In 1984, we described GABA receptors in rat cerebellum that were insensitive to both bicuculline and baclofen that became known as GABA_C receptors (Drew *et al.*, 1984; Chebib and Johnston, 2000). Subsequent molecular biological and neuropharmacological studies showed that GABA_C receptors were homomeric ionotropic GABA receptors made up of ρ -subunits, that had been first cloned from the retina, and having a distinct pharmacological agonist and antagonist profile (Bormann and Feigenspan, 1995; Johnston, 1996). More recently, the International Union of Basic and Clinical Pharmacology recommended that GABA_C receptors were a subtype of GABA_A

receptors and should be classified as GABA_A receptors (Olsen and Sieghart, 2008, see also Alexander *et al.*, 2011). It was 'especially recommended that the name GABA_C receptor should not be used as the sole name for the ρ -receptors in an article including, especially, the title and abstract'. This recommendation has not met with general acceptance with investigators actually working on GABA_C receptors (including me), particularly those working in the retina, where there is a very interesting interaction between GABA_A and GABA_C receptors (Lukasiewicz and Shields, 1998; Bormann and Feigenspan, 2001). Based on data in SciFinder, more than 80 publications in major journals including the British Journal of Pharmacology (Gasulla *et al.*, 2012) since 2009 have contained the term GABA_C in their titles or abstracts.

Bicuculline-sensitive GABA receptors are part of the superfamily of Cys-loop pentameric ligand-gated ion channel receptors that include nicotinic acetylcholine, glycine and 5HT₃ receptors (Chebib and Johnston, 2000). A variety of GABA_A protein subunits have been described including α 1–6, β 1–3, γ 1–3, δ , ϵ , θ and π that can combine to provide a large multiplicity of GABA_A receptor subtypes with a diversity of function and pharmacology (Olsen and Sieghart, 2009). The combinatorial capability of heteromeric ligand-gated ion channels to afford a large variety of functional receptors by different combinations of a relatively small number of protein subunits rivals the wide variety of G-protein coupled receptors produced from a large number of protein subunits widely quoted as being the major drug targets for CNS drug development. Both ligand-gated and G-protein coupled receptors offer a seemingly bewildering array of drug targets.

Functional studies on recombinant and native GABA_A receptors have indicated a lack of dependence on subunit composition for the antagonist action of bicuculline (Krishek *et al.*, 1996). GABA_A receptor subtypes that contain the α 6 subunit are less sensitive to bicuculline, while GABA_C receptors containing ρ subunits are insensitive to bicuculline (Ng *et al.*, 2011).

While the potency of agonists and modulators of GABA_A receptors varies with subunit composition, the potency of

most antagonists is largely independent of receptors subunit composition (Ebert *et al.*, 1997). Whereas, antagonists show comparable potencies in binding and functional studies, the potency of agonists in binding studies are generally two to three orders of magnitude higher than the agonist potencies measured electrophysiologically. These findings indicate that the GABA_A receptor mechanisms transducing binding into physiological response, rather than the binding *per se*, is more dependent on the receptor subunit composition (Ebert *et al.*, 1997). Studies on the differential sensitivity of three inbred strain of mice to bicuculline suggest that genetic factors influence the convulsant action of bicuculline (Freund *et al.*, 1987).

How does bicuculline act on GABA_A receptors?

Bicuculline acts as a competitive antagonist at GABA_A receptors in that it competitively inhibits GABA binding to these receptors and, in turn, GABA competitively inhibits bicuculline binding (Andrews and Johnston, 1979). This evidence is supported by functional studies. Single channel studies have shown that bicuculline reduces GABA activated conductance by reducing both channel open times and opening frequency (Macdonald *et al.*, 1989). Studies indicate that bicuculline binds at the orthosteric site, interacting with additional sites to GABA resulting in stabilization of the receptor in a closed state and in this sense bicuculline has been described as an allosteric inhibitor (Ueno *et al.*, 1997). Given that bicuculline is some three times the molecular size of GABA, it is not unexpected that bicuculline could bind to sites to which GABA could not reach outside the orthosteric site.

Key aspects of structure-activity studies on bicuculline as a GABA_A antagonist are summarized in Figure 4 (Johnston, 1991). The cationic nitrogen and the lactone/ester function are important for interaction with the orthosteric site, while the two aromatic rings with methylenedioxy or methoxy




Figure 4

Summary of structure-activity studies on bicuculline and related compounds showing important aspects that contribute to the GABA receptors antagonist properties.

groups are considered important for interaction with the additional sites that contribute to antagonist action.

Why does bicuculline lack activity at some ionotropic GABA receptors?

Zhang and colleagues have studied the structural determinants for bicuculline antagonist activity using individual or combined mutations of nine residues in $\rho 1$ GABA_C and $\alpha 1\beta 2\gamma 2$ GABA_A recombinant receptors (Zhang *et al.*, 2008). They showed that the bicuculline insensitivity of the $\rho 1$ GABA_C receptor was mainly determined by Tyr106, Phe138 and Phe240 residues. They also studied gabazine insensitivity that was highly dependent on Tyr102, Tyr106 and Phe138, consistent with differences between bicuculline and gabazine as GABA_A receptor competitive antagonists. Mutation of Tyr157 to Ser in the $\beta 2$ subunit of $\alpha 1\beta 2\gamma 2$ L rat GABA_A recombinant receptors resulted in bicuculline acting as a weak agonist (Ueno *et al.*, 1997). Mutations in the $\alpha 1$ subunit of $\alpha 1\beta 2\gamma 2$ GABA_A recombinant receptors showed that the substitution in the N-terminal region of the extracellular domain of Phe64 by Leu decreased the affinities of bicuculline methiodide and gabazine by 60- to 200-fold, suggesting 'close functional and structural association of the α -subunit with the agonist/antagonist binding site' (Sigel *et al.*, 1992).

Other actions of bicuculline

Bicuculline and more particularly its quaternary salts have significant actions other than as GABA_A receptors antagonists (Seutin and Johnson, 1999). While the actions of these agents on nicotinic acetylcholine receptors can be readily distinguished from their effects on GABA_A receptors, they reflect the structural similarities between nicotinic and GABA_A receptors, for example, (+)-tubocurarine acts on both but is more potent on nicotinic receptors (Hill *et al.*, 1973). Bicuculline quaternary salts and (+)-tubocurarine also act on SK (small conductance) calcium-activated potassium channels. Detailed studies on the antagonism of recombinant nicotinic acetylcholine receptors by bicuculline led to the suggestion that 'bicuculline may serve as a lead molecule to design new anticholinergic substances' (Demuro *et al.*, 2001). Bicuculline quaternary salts also act as inhibitors of acetylcholinesterase (Breuker and Johnston, 1975). GABA-metabolising enzymes are not influenced by bicuculline (Beart and Johnston, 1972).

Picrotoxin and picrotoxinin

Reports in the late 1950s that picrotoxin blocked the action of GABA at invertebrate inhibitory synapses and the observations in the early 1960s by Eccles and his colleagues (1963) of the reduction of the presynaptic inhibition of monosynaptic reflexes by this convulsant suggested that GABA was a transmitter in the mammalian spinal cord. We experienced difficulties in demonstrating that picrotoxin or its active component picrotoxinin (Figure 5, picrotoxin is a 1:1

mixture of picrotoxinin and relatively inactive picrotin) influenced the effect of GABA on spinal neurones using microelectrophoresis. Because picrotoxinin is poorly soluble and not ionized in water, local concentrations achieved by attempted microelectrophoretic administration were presumably too low. We were unable to reliably repeat the demonstration by others (Davidoff and Aprison, 1968) that microelectrophoretically administered picrotoxin consistently blocked the action of glycine on spinal interneurons (Curtis *et al.*, 1969). Frustration and confusion over our results regarding picrotoxinin led us to search for agents that could reliably distinguish between the effects of GABA and glycine such as we had already demonstrated with strychnine (Curtis *et al.*, 1967).

Using recombinant receptors, bath application of picrotoxin was subsequently shown to be a mixed/non-competitive GABA_A receptor antagonist, in contrast, to the competitive antagonist action of bicuculline (Krishek *et al.*, 1996). In addition to its action as a GABA_A receptor antagonist, picrotoxinin is now known to act upon GABA_C, glycine and 5HT₃ receptors. Picrotoxin is considered to bind in the pore region of these ligand-gated ion channels (Thompson *et al.*, 2010).


Other GABA_A receptor antagonists

A wide variety of GABA_A receptor antagonists now exists. Four of the more interesting antagonists are discussed briefly below (Figure 5). Of these, gabazine is the most widely studied.

Gabazine, as SR 95531, was introduced as a specific, potent and competitive GABA_A antagonist in 1986 as a result of the study of arylaminopyridazine derivatives of GABA (Heaulme *et al.*, 1986). While gabazine and bicuculline have similar actions as antagonists of the activation of GABA_A receptors on hippocampal neurons by GABA, gabazine is much weaker against activation by pentobarbitone and etomidate (Uchida *et al.*, 1996). This and other findings suggests that GABA, pentobarbitone and etomidate act at non-identical sites on these neurons to open chloride ionophores and that these can be distinguished by gabazine and bicuculline (Ueno *et al.*, 1997). As noted above, gabazine and bicuculline interact with different residues on GABA_A receptors (Zhang *et al.*, 2008). Structure-activity studies have been carried out on gabazine, identifying more potent analogues (Iqbal *et al.*, 2011).

Salicylidene salicylhydrazide (SCS) was identified as a potent inhibitor of the activation of GABA_A receptors containing the $\beta 1$ subunit using a high throughput screening assay (Thompson *et al.*, 2004). It produced a maximum inhibition of 56% with an IC₅₀ of 32 nM. The presence of threonine 255 and isoleucine 308 within the $\beta 1$ subunit and the lack of interaction with a range of GABA_A receptor agents, including bicuculline and picrotoxin, suggest that SCS is interacting at a previously unidentified site.

RU5135, a convulsant steroid derivative, is one of the most potent GABA_A antagonists, being some 500 times more potent than bicuculline in binding studies (Hunt and Clements-Jewery, 1981). In electrophysiological studies, it acted as a competitive antagonist with a pA₂ of 8.31 and

**Figure 5**

Structures of some GABA_A receptors antagonists unrelated to bicuculline.

appeared to share a common site of action with bicuculline (Simmonds and Turner, 1985). However, it lacks specificity, as it is also a glycine antagonist sharing a common site of action with strychnine.

4-(3-Biphenyl)-5-(4-piperidyl)-3-isoxazolol (3-biphenyl-4-PIOL, 9 K in Frølund *et al.*, 2007) was derived from 4-PIOL, a low efficacy partial GABA_A receptor agonist. It is a selective GABA_A receptor antagonist acting in the low nM range. Studies on analogues of 4-PIOL substituted in the four-position of the isoxazole ring yielded GABA_A receptor antagonists of increased potency, and there was a linear correlation between the lipophilicity of the four-substituent and antagonist activity, providing evidence for a hydrophobic binding pocket at the GABA recognition site (Frølund *et al.*, 2007).

Not all GABA_A receptor antagonists are convulsants

The basic tenet for the successful search for bicuculline as a GABA receptor antagonist was that such an agent would be a convulsant on systemic administration, just as strychnine was a convulsant acting on glycine receptors. We and others have described a series of terpenoids structurally related to picrotoxinin, including bilobalide (Figure 4), from *Ginkgo*

biloba that are relatively potent antagonists at ionotropic GABA receptors (Sasaki *et al.*, 1999b; Huang *et al.*, 2003; Ivic *et al.*, 2003). These agents also antagonize glycine and 5HT₃ receptors (Hawthorne *et al.*, 2006; Thompson *et al.*, 2011), but they are not convulsants on systemic administration to mammals. Indeed, they act as anticonvulsants (Sasaki *et al.*, 1999a) and neuroprotectants (DeFeudis, 2002; Huang *et al.*, 2012). They appear to have diverse and competing actions on CNS neurotransmission including reducing the release of L-glutamate thus, reducing excitation (Johns *et al.*, 2002) and inhibiting GABA synthesis via inhibition of glutamate decarboxylase (Sasaki *et al.*, 1999a). The lack of convulsant actions of these terpenoids may be masked by their multiplicity of actions on a variety of neurotransmitters.

Advantages of an antagonist

Thus bicuculline became a useful tool for probing GABA-mediated synaptic inhibition. The subsequent discovery of bicuculline-insensitive GABA receptors, its instability and actions of bicuculline not related to GABA receptors meant that caution had to be taken in interpreting results using bicuculline.

Until the discovery of bicuculline as a GABA receptor antagonist, along with many scientists, David Curtis did not believe that GABA was a neurotransmitter in the spinal cord.

This was based on the relative ubiquity of GABA's action as a neuronal depressant in the brain and spinal cord, and the difficulties in showing that its action could be blocked by picrotoxin. He writes in his autobiography (Curtis, 2006) about his participation in the May 1959 symposium 'Inhibition in the Nervous System and Gamma-Aminobutyric Acid' organized by Gene Roberts in Duarte, California: 'My paper dealt with the effects of GABA, L-GLUT, and related amino acids on spinal neurons, and my negative conclusions related to transmitter functions were unfortunately based on a faulty technique and incorrect assumptions.' Nonetheless, Curtis continued to investigate the function of GABA in the CNS. Spurred on by the discovery of strychnine as a glycine antagonist (Curtis *et al.*, 1967), Curtis actively encouraged what turned out to be a successful search for an equivalent GABA antagonist and with great enthusiasm demonstrated that this GABA antagonist was able to reduce the strychnine-insensitive postsynaptic inhibitions of Deiters' cells, Purkinje cells, pyramidal cells in the cerebral and hippocampal cortices and thalamocortical relay cells. Such studies provided substantive evidence for the role of GABA as an inhibitory neurotransmitter in the CNS.

David Curtis became a convert to the concept of GABA as an inhibitory neurotransmitter as a result of the breakthrough in the use of bicuculline as a GABA antagonist. This is reminiscent of his great mentor, Sir John Eccles, becoming a convert to the concept of chemical neurotransmission as a result of significant technological improvements in electrophysiological recordings. Interestingly, just as it is now known that electrical communication between neurones can take place, we now also know that GABA has many other functions including acting as a trophic factor to influence events such as proliferation, migration, differentiation, synapse maturation and cell death (Owens and Kriegstein, 2002). In mammals, GABA is found in many organs outside of the CNS where it serves various functions. GABA is involved in cell proliferation and migration, and may play a role in cancer. Recent evidence implicates GABA receptors in mucus overproduction in asthma acting on airway epithelial cells. GABA regulates insulin secretion from pancreatic β cells in concert with changes in glucose concentration and may be involved with type 1 diabetes (Braun *et al.*, 2010). Functional GABA receptors have also been described in T cells and macrophages (Tian *et al.*, 1999; Shiratsuchi *et al.*, 2009). Thus, in addition to neurotransmission in the CNS, GABA is involved in asthma, cancer, diabetes and the immune system (Hanrahan and Johnston, 2009). The discovery of agents (not only antagonists but also the full range of allosteric, full and partial agonists, and negative, neutralizing and positive modulators) that selectively influence the function of GABA in these systems remains a challenge for medicinal chemists and pharmacologists, since such molecules will have wide applicability in diverse clinical conditions.

Acknowledgements

The author is indebted to the many colleagues and students that contributed to the studies on bicuculline and related substances. He is grateful to Philip Beart, Jane Hanrahan and

Ken Mewett for their constructive input into the writing of this review.

Conflict of interest

The author has no conflict of interest.

References

- Alexander SPH, Mathie A, Peters JA (2011). Guide to Receptors and Channels (GRAC), 5th edn. Br J Pharmacol 164 (Suppl. 1): S1–S324.
- Andrews PR, Johnston GAR (1979). GABA agonists and antagonists. Biochem Pharmacol 28: 2697–2702.
- Beart PM, Johnston GAR (1972). Bicuculline and GABA-metabolising enzymes. Brain Res 38: 226–227.
- Bormann J, Feigenspan A (1995). GABA_C receptors. Trends Neurosci 18: 515–519.
- Bormann J, Feigenspan A (2001). GABA_C receptors: structure, function and pharmacology. Handb Exp Pharmacol 150: 271–296.
- Braun M, Ramracheya R, Bengtsson M, Clark A, Walker JN, Johnson PR *et al.* (2010). γ -aminobutyric acid (GABA) is an autocrine excitatory transmitter in human pancreatic beta-cells. Diabetes 59: 1694–1701.
- Breuker E, Johnston GAR (1975). Inhibition of acetylcholinesterase by bicuculline and related alkaloids. J Neurochem 25: 903–904.
- Chebib M, Johnston GAR (2000). GABA-activated ligand gated ion channels: medicinal chemistry and molecular biology. J Med Chem 43: 1427–1447.
- Curtis DR (2006). David R. Curtis Autobiography. Hist Neurosci Autobiogr 5: 171–225.
- Curtis DR, Hosli L, Johnston GAR (1967). Inhibition of spinal neurons by glycine. Nature 215: 1502–1503.
- Curtis DR, Duggan AW, Johnston GAR (1969). Glycine, strychnine, picrotoxin and spinal inhibition. Brain Res 14: 759–762.
- Curtis DR, Duggan AW, Felix D, Johnston GAR (1970a). Bicuculline and central GABA receptors. Nature 228: 676–677.
- Curtis DR, Duggan AW, Felix D, Johnston GAR (1970b). GABA, bicuculline and central inhibition. Nature 226: 1222–1224.
- Curtis DR, Duggan AW, Felix D, Johnston GAR (1971). Bicuculline, an antagonist of GABA and synaptic inhibition in the spinal cord of the cat. Brain Res 32: 69–96.
- Davidoff RA, Aprison MH (1968). Picrotoxin antagonism of the inhibition of interneurons by glycine. Life Sci 8: 107–112.
- DeFeudis FV (2002). Bilobalide and neuroprotection. Pharmacol Res 46: 565–568.
- Demuro A, Palma E, Eusebi F, Miledi R (2001). Inhibition of nicotinic acetylcholine receptors by bicuculline. Neuropharmacology 41: 854–861.
- Drew CA, Johnston GAR, Weatherby RP (1984). Bicuculline-insensitive GABA receptors: studies on the binding of (-)-baclofen to rat cerebellar membranes. Neurosci Lett 52: 317–321.
- Ebert B, Thompson SA, Saounatsou K, McKernan R, Krosgaard-Larsen P, Wafford KA (1997). Differences in

- agonist/antagonist binding affinity and receptor transduction using recombinant human γ -aminobutyric acid type A receptors. *Mol Pharmacol* 52: 1150–1156.
- Eccles JC, Schmidt R, Willis RD (1963). Pharmacological studies on presynaptic inhibition. *J Physiol* 168: 500–530.
- Editorial (1970). Advantages of an antagonist. *Nature* 226: 1199–1200.
- Freund RK, Marley RJ, Wehnder JM (1987). Differential sensitivity to bicuculline in three inbred mouse strains. *Brain Res Bull* 18: 657–662.
- Frølund B, Jensen LS, Storustovu SI, Stensbol TB, Ebert B, Kehler J *et al.* (2007). 4-aryl-5-(4-piperidyl)-3-isoxazolol GABA_A antagonists: synthesis, pharmacology, and structure-activity relationships. *J Med Chem* 50: 1988–1992.
- Gasulla J, Gonzalez ANB, Calvo DJ (2012). Nitric oxide potentiation of the homomeric $\rho 1$ GABA_C receptor function. *Br J Pharmacol* 167: 1369–1377.
- Godfraind JM, Krnjevic K, Pumain R (1970). Doubtful value of bicuculline as a specific antagonist of GABA. *Nature* 228: 675–676.
- Hanrahan JR, Johnston GAR (2009). Synthesis of γ -aminobutyric acid analogues. In: Hughes AB (ed.). *Amino Acids, Peptides and Proteins in Organic Chemistry*. Vol. Volume 1, Origins and synthesis of amino acids. Wiley VCH: Weinheim, pp. 573–689.
- Hawthorne R, Cromer BA, Ng HL, Parker MW, Lynch JW (2006). Molecular determinants of ginkgolide binding in the glycine receptor pore. *J Neurochem* 98: 395–407.
- Heaulme M, Chambion JP, Leyris R, Molimard JC, Wermuth CG, Bizuere K (1986). Biochemical characterization of the interaction of three pridaziny-GABA derivatives with the GABA_A receptor site. *Brain Res* 384: 224–231.
- Hill DR, Bowery NG (1981). ³H-Baclofen and ³H-GABA bind to bicuculline insensitive GABA sites in rat brain. *Nature* 290: 149–152.
- Hill RG, Simmonds MA, Straughan DW (1973). A comparative study of some convulsant substances as γ -aminobutyric acid antagonists in the feline cerebral cortex. *Br J Pharmacol* 49: 37–51.
- Hoffman M (1994). Lost in the cultural revolution. *Am Sci* 82: 18–19.
- Huang M, Qian Y, Guan T, Huang L, Tang X, Li Y (2012). Different neuroprotective responses of Ginkgolide B and bilobalide, the two Ginkgo components, in ischemic rats with hyperglycemia. *Eur J Pharmacol* 677: 71–76.
- Huang SH, Duke RK, Chebib M, Sasaki K, Wada K, Johnston GAR (2003). Bilobalide, a sesquiterpene trilactone from *Ginkgo biloba*, is an antagonist at recombinant $\alpha 1, \beta 2, \gamma 2L$ GABA_A receptors. *Eur J Pharmacol* 464: 1–8.
- Hunt P, Clements-Jewery S (1981). A steroid derivative, R 5135, antagonizes the GABA/benzodiazepine receptor interaction. *Neuropharmacology* 20: 357–361.
- Iqbal F, Ellwood R, Mortensen M, Smart TG, Baker JR (2011). Synthesis and evaluation of highly potent GABA_A receptor antagonists based on gabazine (SR-95531). *Bioorg Med Chem Lett* 21: 4252–4254.
- Ivic L, Sands TT, Fishkin N, Nakanishi K, Kriegstein AR, Stromgaard K (2003). Terpene trilactones from *Ginkgo biloba* are antagonists of cortical glycine and GABA_A receptors. *J Biol Chem* 278: 49279–49285.
- Johns L, Sinclair AJ, Davies JA (2002). Effects of bilobalide on hypoxia/hypoglycemia-stimulated glutamate efflux from rat cortical brain slices. *Neurochem Res* 27: 369–371.
- Johnston GAR (1985). First Sino-Australian workshop in pharmacology. *Trends Pharmacol Sci* 159.
- Johnston GAR (1991). GABA_A antagonists. *Semin Neurosci* 3: 205–210.
- Johnston GAR (1996). GABA_C receptors: relatively simple transmitter-gated ion channels? *Trends Pharmacol Sci* 17: 319–323.
- Johnston GAR, Beart PM, Curtis DR, Game CJ, McCulloch RM, MacIachlan RM (1972). Bicuculline methochloride as a GABA antagonist. *Nat New Biol* 240: 219–220.
- Krishek BJ, Moss SJ, Smart TG (1996). A functional comparison of the antagonists bicuculline and picrotoxin at recombinant GABA_A receptors. *Neuropharmacology* 35: 1289–1298.
- Lukasiewicz PD, Shields CR (1998). Different combinations of GABA_A and GABA_C receptors confer distinct temporal properties to retinal synaptic responses. *J Neurophysiol* 79: 3157–3167.
- Macdonald RL, Rogers CJ, Twyman ER (1989). Kinetic properties of the GABA_A receptor main conductance state of mouse spinal cord neurones in culture. *J Physiol (Lond)* 419: 479–499.
- McLennan H (1970). Bicuculline and inhibition of crayfish stretch receptor neurones. *Nature* 228: 674–675.
- Ng CK, Kim HL, Gavande N, Yamamoto I, Kumar RJ, Mewett KN *et al.* (2011). Medicinal chemistry of ρ GABA_C receptors. *Future Med Chem* 3: 197–209.
- Olsen RW, Sieghart W (2008). International Union of Pharmacology. LXX. Subtypes of γ -aminobutyric acid (A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev* 60: 243–260.
- Olsen RW, Sieghart W (2009). GABA A receptors: subtypes provide diversity of function and pharmacology. *Neuropharmacology* 56: 141–148.
- Olsen RW, Ban M, Miller T, Johnston GAR (1975). Chemical instability of the GABA antagonist bicuculline under physiological conditions. *Brain Res* 98: 383–387.
- Owens DF, Kriegstein AR (2002). Is there more to GABA than synaptic inhibition? *Nat Rev Neurosci* 3: 715–727.
- Pong SF, Graham LT (1972). N-methyl bicuculline, a convulsant more potent than bicuculline. *Brain Res* 42: 486–490.
- Roberts E, Kuriyama K, Haber B (1970). Biochemistry of synaptic inhibition at the cellular level: the GABA system. *Adv Biochem Psychopharmacol* 2: 139–161.
- Sasaki K, Hatta S, Haga M, Ohshika H (1999a). Effects of bilobalide on γ -aminobutyric acid levels and glutamic acid decarboxylase in mouse brain. *Eur J Pharmacol* 367: 165–173.
- Sasaki K, Oota I, Wada K, Inomata K, Ohshika H, Haga M (1999b). Effects of bilobalide, a sesquiterpene in *Ginkgo biloba* leaves, on population spikes in rat hippocampal slices. *Comp Biochem Physiol C Comp Pharmacol Toxicol* 124: 315–321.
- Seutin V, Johnson SW (1999). Recent advances in the pharmacology of quaternary salts of bicuculline. *Trends Pharmacol Sci* 20: 268–270.
- Shiratsuchi H, Kouatli Y, Yu GX, Marsh HM, Basson MD (2009). Propofol inhibits pressure-stimulated macrophage phagocytosis via the GABA_A receptor and dysregulation of p130cas phosphorylation. *Am J Physiol Cell Physiol* 296: C1400–C1410.

- Sigel E, Baur R, Kellenberger S, Malherbe P (1992). Point mutations affecting antagonist affinity and agonist dependent gating of GABA receptor channels. *EMBO J* 11: 2017–2023.
- Simmonds MA, Turner JP (1985). Antagonism of inhibitory amino acids by the steroid derivative RU5135. *Br J Pharmacol* 84: 631–635.
- Thompson AJ, Lester HA, Lummis SC (2010). The structural basis of function in Cys-loop receptors. *Q Rev Biophys* 43: 449–499.
- Thompson AJ, Jarvis GE, Duke RK, Johnston GA, Lummis SC (2011). Ginkgolide B and bilobalide block the pore of the 5-HT₃ receptor at a location that overlaps the picrotoxin binding site. *Neuropharmacology* 60: 488–495.
- Thompson SA, Wheat L, Brown NA, Wingrove PB, Pillai GV, Whiting PJ *et al.* (2004). Salicylidene salicylhydrazide, a selective inhibitor of β 1-containing GABA_A receptors. *Br J Pharmacol* 142: 97–106.
- Tian J, Chau C, Hales TG, Kaufman DL (1999). GABAA receptors mediate inhibition of T cell responses. *J Neuroimmunol* 96: 21–28.
- Tsou K (1987). Mechanism of morphine analgesia and hybridization approaches to study the role of opioid peptides in acupuncture. In: Rand MJ, Raper C (eds). *Pharmacology – Proceedings of the Xth International Congress of Pharmacology*, Edn. Elsevier: Sydney, pp. 225–234.
- Uchida I, Cestari IN, Yang J (1996). The differential antagonism by bicuculline and SR95531 of pentobarbitone-induced currents in cultured hippocampal neurons. *Eur J Pharmacol* 307: 89–96.
- Ueno S, Bracamontes J, Zorumski C, Weiss DS, Steinbach JH (1997). Bicuculline and gabazine are allosteric inhibitors of channel opening of the GABA_A receptor. *J Neurosci* 17: 625–634.
- Welch AD, Henderson VE (1934). A note on bicucine. *J Pharmacol Exp Ther* 51: 492–494.
- Werman R, Davidoff RA, Aprison MH (1967). Inhibition of motoneurons by iontophoresis of glycine. *Nature* 214: 581–583.
- Zhang J, Xue F, Chang Y (2008). Structural determinants for antagonist pharmacology that distinguish the ρ 1 GABA_C receptor from GABA_A receptors. *Mol Pharmacol* 74: 941–951.